

Effect of fertilizers for *Phaseolus vulgaris* L. productivity in Western Forest-Steppe of Ukraine

I. Didur¹, O. Chynchyk², H. Pantsyreva^{1*} , S. Olifirovych², V. Olifirovych³, O. Tkachuk²

¹Vinnitsia National Agrarian University, Vinnitsia, Ukraine

²State Agrarian and Engineering University in Podilia, Kamyanets-Podilsky, Ukraine

³Bukovinian State Agricultural Research Station of the Institute of Agriculture of the Carpathian Region NAAS

*Corresponding author Email: pantsyreva@vsau.vin.ua

Received: 05.01.2021. Accepted: 05.02.2021

The scientific article summarizes the improving the efficiency of growing legumes by selecting varieties, primary tillage mode, and optimization of plant nutrition in the Western Forest-Steppe. Studies have shown that legumes are the most demanding crops. In the Western Forest-Steppe, the Bukovynka variety of legumes became more productive when inoculating seeds with Rhizobophyte, applying mineral fertilizer (N30P60K60), foliar fertilization of crystallin, the grain yield of legumes was 2.69 t/ha. A high correlation is observed between the number of beans and the number of grains per plant ($r = 0.98$), the mass of grains from one plant and the mass of 1000 grains ($r = 0.97$), number of grains per 1 plant, and the mass of grains per plant ($r = 0.96$). The moderate correlation was between the number of grains in the bean ($r = 0.63$) and the weight of 1000 grains ($r = 0.66$). A low correlation ($r = 0.45$) was observed between the number of beans per plant and the bean's number of grains. Negative correlations we registered for the number of beans per plant and a mass of 1000 grains ($r = -0.37$) and the number of grains per one plant ($r = -0.17$). Thus, we revealed the Bukovynka variety provided the highest level of legume grain yield during inoculation of seeds with Rhizobophyte, applying mineral fertilizer at the dose of N30P60K60, and two foliar fertilization of crops with Kristalon.

Keywords: *Phaseolus vulgaris* L., productivity, variety, mineral fertilizers, foliar fertilization, grain yield.

Introduction

During the reform of the agro-industrial complex of Ukraine and animal production reduction, high-protein crop products' production became important. As a result, the demand for legume seeds has risen sharply in recent years. Legumes' particular importance in solving humankind's protein problem is determined primarily by the high protein content, a significant number of essential and critical amino acids, high solubility, and nutritional value.

Legumes contain an average of 24% protein, which in the amino acid composition is close to proteins of animal origin. We know these plants enrich the soil with valuable organic matter, nitrogen, replenish the arable layer with phosphorus, potassium, calcium, improve soil structure, and increase fertility. They are the best precursors for most crop rotation and the most valuable green manure; we can grow them without using nitrogen fertilizers, which account for up to 30% of energy consumption in intensive technologies. We can consider that the utilization rate of nitrogen from mineral fertilizers is only 50%, a significant part of them contaminates groundwater with nitrates, and living organisms entirely use biological nitrogen.

Legumes are crops that requires high soil fertility. About 90-95% of P, K, and Ca is absorbed in a brief period – only 50-65 days from the moment of emergence. Legumes are the most demanding of all legumes. According to Stakanov F., legumes responded well to nitrogen fertilizers at a dose of N45 – yields increased by 11%. When growing legumes on chernozems, we need to use phosphorus and phosphorus and potassium fertilizers. After all, soil acidity and low phosphorus levels are critical limiting factors for increasing bean yields.

It is also essential to choose an effective bean inoculant. According to American scientists (Gepts et al., 1988), only six Rhizobium strains out of twelve were effective for *Phaseolus vulgaris* L. Thus, treatment of seeds with Glomus® and Rhizotorphin® increased the grain yield of legumes by 2.3-4.5 kg/ha. According to D. Shlyakhturova, the combination in the fertilizer system of ordinary nitrogen and mineral fertilizers created the best conditions for forming this culture's plant productivity. It is established that on the dark gray podzolic soil in the Lviv region conditions, the highest yield of legumes was obtained by applying mineral fertilizers in the pre-sowing cultivation in the dose N30P60K60. In the conditions of the Northern Forest-Steppe of Ukraine, the maximum yield of legumes was obtained by inoculating seeds and applying N60P60K60 (Bakhmat,2016) In Bulgaria, the yield of legumes increased in all variants of the experiment with mineral

fertilizers compared to the control (Oliveira et al., 2000). In Brazil (Echeverrigaray et al., 1993), phosphorus and molybdenum's introduction has increased the yield of legumes, especially when inoculating seeds with nodule bacteria.

Materials and methods

Scientific research was performed by conducting field and laboratory experiments. The research was conducted following generally accepted methods. The soil of the experimental field is leached black soil, low humus, on carbonate loam. The humus content in the soil layer is 0-30 cm (according to Tyurin is 3.86-4.11%; easily hydrolyzing nitrogen (according to Cornfield) – 111-121 mg/kg, mobile phosphorus and metabolic potassium (according to Chirikov) – 90 and 179 mg/kg of soil, respectively. The absorption capacity and the number of absorbed bases vary, respectively, between 33-36 and 30-33 mg-eq/100 g of soil. Hydrolytic acidity is 0.76-0.87 mg-eq/100 g soil, the degree of saturation of the basics – 94.7-99.0%. The solid phase density is 2.58 g/cm³, the density of the soil structure is 1.14-1.25 g/cm³, the total porosity is 52-59%. The maximum soil hygroscopicity is 5.2%; the lowest moisture content is 23.4%, the total field capacity is 41.2%.

The climate of the southwestern part of the Forest-Steppe of Ukraine is warm, with sufficient moisture. The average radiation balance in the region for the year is 43.3 kcal/cm², and for the growing season of sugar sorghum – 137.73 kJ/cm². Most of PAR come in June and July. From May to September, 3/4 of the annual amount of heat comes to the soil surface. Annual precipitation ranges from 550-700 mm, 3/4 of which falls during the warm season. The hydrothermal coefficient in the region is 1.4. The weather conditions of the sugar sorghum vegetation period in 2017-2019 years had the following features: with the average long-term rainfall and the sum of temperatures respectively 345 mm and 2903 °C, during the years these indicators fluctuated within such limits.

During the research period, two experiments were conducted. The study examined the effect of fertilizers on grain yield of common bean varieties. Varieties of common beans, entered in the State Register of varieties allowed for distribution in Ukraine Nadiya, Bukovynka.

Results and discussion

Table 1. Grain productivity of bean varieties, depending on power supply systems, t/ha.

| Fertilizers (factor A) | Yield by year | | | | | average |
|--|---------------|------|------|------|------|---------|
| | 2015 | 2016 | 2017 | 2018 | 2019 | |
| grade (factor B) – Nadiya | | | | | | |
| No fertilizers (c.) | 2.24 | 1.71 | 2.16 | 2.14 | 2.07 | 2.07 |
| P ₆₀ K ₆₀ | 2.35 | 2.09 | 2.27 | 2.25 | 2.24 | 2.24 |
| N ₃₀ P ₆₀ K ₆₀ | 2.46 | 2.12 | 2.33 | 2.29 | 2.30 | 2.30 |
| Rhizobophyte | 2.38 | 1.76 | 2.23 | 2.31 | 2.17 | 2.17 |
| P ₆₀ K ₆₀ + Rhizobophyte | 2.44 | 2.02 | 2.42 | 2.35 | 2.31 | 2.31 |
| N ₃₀ P ₆₀ K ₆₀ + Rhizobophyte | 2.46 | 2.12 | 2.43 | 2.36 | 2.32 | 2.34 |
| Cristalon | 2.36 | 1.99 | 2.28 | 2.33 | 2.24 | 2.24 |
| P ₆₀ K ₆₀ + Cristalon | 2.57 | 2.11 | 2.25 | 2.39 | 2.33 | 2.33 |
| N ₃₀ P ₆₀ K ₆₀ + Cristalon | 2.59 | 2.24 | 2.37 | 2.44 | 2.41 | 2.41 |
| Rhizobophyte + Cristalon | 2.42 | 1.83 | 2.35 | 2.40 | 2.25 | 2.25 |
| P ₆₀ K ₆₀ + Rhizobophyte + Cristalon | 2.51 | 2.19 | 2.38 | 2.48 | 2.39 | 2.39 |
| N ₃₀ P ₆₀ K ₆₀ + Rhizobophyte + Cristalon | 2.64 | 2.32 | 2.42 | 2.54 | 2.48 | 2.48 |
| grade (factor B) – Bukovynka | | | | | | |
| No fertilizers | 2.57 | 1.82 | 2.43 | 2.50 | 2.33 | 2.33 |
| P ₆₀ K ₆₀ | 2.77 | 1.96 | 2.52 | 2.71 | 2.49 | 2.49 |
| N ₃₀ P ₆₀ K ₆₀ | 2.84 | 2.06 | 2.56 | 2.82 | 2.62 | 2.57 |
| Rhizobophyte | 2.65 | 1.86 | 2.47 | 2.54 | 2.38 | 2.38 |
| P ₆₀ K ₆₀ + Rhizobophyte | 2.81 | 1.95 | 2.54 | 2.74 | 2.51 | 2.51 |
| N ₃₀ P ₆₀ K ₆₀ + Rhizobophyte | 2.87 | 2.08 | 2.62 | 2.83 | 2.60 | 2.60 |
| Cristalon | 2.81 | 1.91 | 2.68 | 2.76 | 2.54 | 2.54 |
| P ₆₀ K ₆₀ + Cristalon | 2.91 | 2.09 | 2.75 | 2.89 | 2.66 | 2.66 |
| N ₃₀ P ₆₀ K ₆₀ + Cristalon | 2.92 | 2.13 | 2.78 | 2.89 | 2.68 | 2.68 |
| Rhizobophyte + Cristalon | 2.84 | 1.90 | 2.71 | 2.79 | 2.56 | 2.56 |
| P ₆₀ K ₆₀ + Rhizobophyte + Cristalon | 2.93 | 2.12 | 2.73 | 2.90 | 2.67 | 2.67 |
| N ₃₀ P ₆₀ K ₆₀ + Rhizobophyte + Cristalon | 2.93 | 2.15 | 2.78 | 2.91 | 2.69 | 2.69 |

HIP₀₅: 2015 A – 0.07; B – 0.09; AB – 0.19; 2016 A – 0.09; B – 0.10; AB – 0.23; 2017 A – 0.09; B – 0.10; AB – 0.20; 2018 A – 0.10; B – 0.12; AB – 0.22; 2019 A – 0.08; B – 0.07; AB – 0.14; average A – 0.07; B – 0.03; AB – 0.09;

(c.) – control.

We found that the average yield of legumes under control in 2015-2019 (Nadiya variety without fertilizer) was 2.07 t/ha. Among the studied varieties in the variant without the use of fertilizers, the highest grain yield was in the variety of legumes Bukovynka - 2.33 t/ha. Our application of fertilizers significantly increased the yield of legumes. Thus, when applying P₆₀K₆₀, the yield fluctuated in the range of 2.24-2.49 t/ha or 8.2-9.4% more than the control. According to the control, the application of N₃₀P₆₀K₆₀ contributed to a further increase in yield to 2.29-2.62 t/h a, which was 11.1-12.4% more. To a lesser extent, legumes' productivity was affected by inoculation of seeds with Rhizobophyte: grain yield of Nadiya variety increased by 0.1 t/ha or 4.8%, Bukovynka variety – by 0.05 t/ha. The combination of inoculation of seeds with Rhizobophyte and application of phosphorus and potassium fertilizers (P₆₀K₆₀) created more optimal conditions for the formation of high productivity of bean varieties. It should be noted that the combination of biological (Rhizobophyte) and mineral (N₃₀) nitrogen on the background of P₆₀K₆₀ contributed to increased grain yield of legumes. Thus, the average grain yield of legumes of the Nadiya variety was 2.31 t/ha, and the Bukovynka variety – 2.51 t/ha. Foliar fertilization of bean crops with Cristalon was also adequate.

In Nadiya variety, treatment with Cristalon provided an increase in yield of 0.17 t/ha, or as much as the application of phosphorus and potassium fertilizers at a dose of P₆₀K₆₀ (Table 1). Against the background of P₆₀K₆₀, the increase in grain yield in this variety was 0.13 t/ha, against the background of N₃₀P₆₀K₆₀ – 0.11 t/ha. The soil nutrition of plants, especially with the application of mineral fertilizer at a dose of N₃₀P₆₀K₆₀, reduced the increase from the use of only Cristalon. With the joint use of Rhizobophyte and Cristalon, the yield increase was 0.18 t/ha, while Rhizobophyte alone was 0.1 t/ha. In our opinion, this is due to the fact that after receiving an additional source of nitrogen nutrition, bean plants spent less nutrients on the functioning of the legume-rhizobial apparatus, and more was spent on the formation of grain yields. Two-factor analysis of variance obtained data showed that the share of the factor "grade" on the yield of legumes was 52%, the factor "fertilizer" – 35%, and the interaction of factors such as "grade" and "fertilizer" – 1%. Other unaccounted for factors were 12%.

The correlation analysis of grain growing indicators of Bukovynka legumes showed a high density of the relationship between grain yield and grain weight per plant ($r = 0.91$), there is also a high level of relationship between yield indicators and the number of grains in legumes ($r = 0.89$), as well as the number of grains per plant ($r = 0.81$). The average relationship between the yield of Bukovynka legumes and the number of legumes per plant is slightly weaker ($r = 0.70$), and the relationship between the value of yields and the weight of 1000 grains is 0.33 (Table 2).

Table 2. Coefficients of correlation between indicators of structural elements and grain yield of Bukovynka legumes (average for 2015-2019).

| Indexes | Number of beans per one plant, pcs. | Number of grains in beans, pcs. | Number of grains per one plant, pcs. | Mass of grains from one plants, g | Mass 1000 grains, g | Yield, t/ha |
|----------------------------------|-------------------------------------|---------------------------------|--------------------------------------|-----------------------------------|---------------------|-------------|
| Number of beans per plant, pcs. | 1.00 | - | - | - | - | - |
| Number of grains in beans, pcs. | 0.45* | 1.00 | - | - | - | - |
| Number of grains per plant, pcs. | 0.98* | 0.63* | 1.00 | - | - | - |
| Mass of grains on one plant, g | 0.89* | 0.81* | 0.96* | 1.00 | - | - |
| Mass 1000 grains, g | -0.37* | 0.66* | -0.17* | 0.97* | 1.00 | - |
| Yield, t/ha | 0.70* | 0.89* | 0.81* | 0.91* | 0.33* | 1.00 |

A high correlation is observed between the number of beans and the number of grains per plant ($r=0.98$), the mass of grains from one plant and the mass of 1000 grains ($r=0.97$), number of grains per 1 plant, and the mass of grains per plant ($r=0.96$). The moderate correlation was between the number of grains in the bean ($r=0.63$) and the weight of 1000 grains ($r=0.66$). A low correlation ($r=0.45$) was observed between the number of beans per plant and the bean's number of grains. Negative correlations we registered for the number of beans per plant and a mass of 1000 grains ($r = -0.37$) and the number of grains per one plant ($r = -0.17$). Thus, we established the Bukovynka variety provided the highest level of legume grain yield during inoculation of seeds with Rhizobophyte, applying mineral fertilizer at the dose of N₃₀P₆₀K₆₀, and two foliar fertilization of crops with Kristalon.

References

- Albinus M. (2008). Effects of land use practices on livelihoods in the transboundary sub-catchments of the Lake Victoria Basin. *African Journal of Environmental Science and Technology*, 2(10), 309-317.
- Bakhmat O.M. (2010). Influence of agrotechnical measures on soybean productivity in the conditions of the western region of Ukraine. *Feed and feed production: interdepartmental. topic. Vinnytsia*, 66, 103-108.
- Bandura V., Mazur V., Yaroshenko L., Rubanenko O. (2019). Research on sunflower seeds drying process in a monolayer tray vibration dryer based on infrared radiation. *INMATEN – Agricultural Engineering*, 57(1), 233-242.
- Begey S. V., Shuvar I. A. (2007). *Ecological Agriculture*. Lviv: Novyi Svit-2000.

- Bulgakov V., Adamchuk V., Kaletnik G., Arak M., Olt J. (2014). Mathematical model of vibration digging up of root crops from soil. *Agronomy Research*, 12(1), 41-58.
- Bulgakov V., Kaletnik H., Goncharuk I., Ivanovs S., Usenko M. (2019). Results of experimental investigations of a flexible active harrow with loosening teeth. *Agronomy Research*, 17(5). 1839–1845.
- Chinchyk O.S. (2012). Influence of fertilizer on yield of legumes in the conditions of the Western Forest-Steppe. Feed and feed production: interdepartmental topic, 72, 64-67.
- Cholovskyi Yu.M. (2010). Osoblyvosti vodospozhyvannia posivamy liupynu vuzkolystoho zalezno vid zastosuvannia mineralnykh dobryv. *Kormy i kormovyrobnytstvo – Forage and feed production*, 66, 146- 147.
- Puyu, V.L., Bakhmat, M.I., Tsvigun, S.A. (2009). Cultural Pasture: Patent No 40618. Podillya State Agrarian Engineering University, UA. Industrial Property, Bulletin No 8.
- Datta, A., Hossain, A., Roy, S. (2019). An Overview on Biofuels and Their Advantages and Disadvantages. *Asian Journal of Chemistry*, 31(8), 1851-1858. DOI: [10.14233/ajchem.2019.22098](https://doi.org/10.14233/ajchem.2019.22098)
- Didur I., Bakhmat M., Chynchyk O., Pansyryeva H., Telekalo N., Tkachuk O. (2020). Substantiation of agroecological factors on soybean agrophytocenoses by analysis of variance of the Right-Bank ForestSteppe in Ukraine, *Ukrainian Journal of Ecology*, 10(5), 54–61.
- Didur I., Pansyryeva H., Telekalo N. (2020). Agroecological rationale of technological methods of growing legumes. *The scientific heritage*, 52, 3–14.
- Didur I.M., Prokopchuk V.M., Pansyryeva H.V. (2019). Investigation of biomorphological and decorative characteristics of ornamental species of the genus *Lupinus* L. *Ukrainian Journal of Ecology*, 9(3), 287-290.
- Echeverrigaray S., Carvalho M.T.V., Pompeu A.S. and Derbyshire E. (1993). Affinity grouping of closely related lines of dry beans (*Phaseolus vulgaris*) through comparative electrophoresis of seed proteins. *Rev. Brasil. Genet*, 16, 759-771.
- Egli D. B. (1999). Variation in leaf starch and sink limitations during seed filling in soybean. *Crop Science*, 39, 1361-1368.
- Environmental Issues of Agriculture. (2010). I.D. Prymak (Ed.). Kyiv: Center of Educational Literature.
- Gepts P., Delgado-Salinas A, Bonet A., Gepts P. (1988). *in Genetic Resources of Phaseolus Beans, The wild relative of Phaseolus vulgaris in Middle America*. Gepts P. (Ed.). Kluwer, Boston, pp 163–184.
- Honcharuk I., Kovalchuk S. (2020). Agricultural Production Greening Management in the Eastern Partnership countries with the EU. Theoretical and practical aspects of the development of the European Research Area. Publishing House Baltija Publishing, Riga, Latvia.H
- Honcharuk I.V., Branitsky Yu.Yu., Tomashuk I.V. (2017). The main aspects of effective formation and use of resource potential in agricultural enterprises (on the example of Vladovo-Lyulynetska DSS IBK and the Central Bank of NAAS of Ukraine). *Economy. Finances. Management: current issues of science and practice*, 10(26), 54-68.
- Kaletnik G.M., Zabolotnyi, G.M., Kozlovskiy S.V. (2011). Innovative models of strategic management economic potential within contemporary economic systems. *Actual Problems of Economics*, 4(118), 11.
- Kaletnik G., Honcharuk, I. (2013). Innovatsiine zabezpechennia rozvytku biopalyvnoi haluzi: svitovyi ta vitchyzniani dosvid [Innovative support for the development of the biofuel industry: world and national experience]. *Biznes Inform [Business Inform]*, 9, 155–160.
- Kaletnik G., Honcharuk I., Okhota Yu. (2020). The Waste-Free Production Development for the Energy Autonomy Formation of Ukrainian Agricultural Enterprises. *Journal of Environmental Management and Tourism*, XI, 3(43), 513-522. DOI:10.14505/jemt.v11.3(43).02
- Kaletnik G. (2018). Production and use of biofuels. Vinnytsia: LLC Nilan-Ltd.
- Kaletnik G., Honcharuk I., Yemchuk T., Okhota Yu. (2020). The World Experience in the Regulation of the Land Circulation. *European Journal of Sustainable Development*, 9(2), 557-568
- Kaletnik G. (2018). Diversification of production of biofuel – as the basis of maintenance of food, power, economic and environmental safety of Ukraine. [Diversification of production of biofuel – as the basis of maintenance of food, power, economic and environmental safety of Ukraine]. *Visnyk aharnoi nauky – Bulletin of agrarian science*, 11, 169-176.
- Kaletnik G.M., Yanovych V.P. (2017). Substantiation of operating and design parameters of a gyration mill for the production of highly active premixes, *Vibrations in engineering and technology*, 84(1), 15-21
- Kaletnik, G., & Lutkovska, S. (2020). Innovative Environmental Strategy for Sustainable Development. *European Journal of Sustainable Development*, 9(2), 89. <https://doi.org/10.14207/ejsd.2020.v9n2p89>
- Kaletnik, G., Shubravskaya, O., Ibatullin, M., Krysanov, D., Starychenko, Y., Tkachenko, K., Varchenko, O. (2019). Features of Food Security of the Country in Conditions of Economic Instability. *Int. J. Manag. Bus. Res*, 9(4), 176-186.
- Kantolic A. G. (2007). Development and seed number in indeterminate soybean as affected by timing and duration of exposure to long photoperiods after flowering. *Annals of Botany*, 99, 925-933.
- Kolesnik, S. (2012). Bacterial fertilizer to optimize nitrogen and phosphorus nutrition soybeans, chickpeas, peas, lentils and commit. *Feed and fodder*, 73, 145-151.
- Kosse, V. Mathew, J. (2017). Design of hammer mills for optimum performance. *Proceeding of the Institution of Mechanical Engineers*, 215, 87-94.
- Kukharchuk V.V., Kaziv S.S., Bykovsky S.A. (2017). Discrete wavelet transformation in spectral analysis of vibration processes at hydropower units, *Przeglad Elektrotechniczny*, 93(5), 65-68.
- Kupchuk I.M., Solona O.V., Derevenko I.A., Tverdokhlib I.V. (2018). Verification of the mathematical model of the energy consumption drive for vibrating disc crusher, *Inmateh – Agricultural Engineering*, 55(2), 111-118.
- Li J., Wang E., Chen W., Chen X. (2008). Genetic diversity and potential for promotion of plant growth detected in nodule endophytic bacteria of soybean grown in Heilongjiang province of China. *Soil Biology & Biochemistry*, 40, 238-246.

- Ma, Z., Wood, C.W., Bransby, D.I. (2001). Impact of row spacing, nitrogen rate, and time on carbon partitioning of switchgrass. *Biomass Bioenergy*, 20, 413–419.
- Malchevskaya, E., Mylenkaya, G. (1981). The comments and Animal Husbandry quality forage analysis. Minsk. Urozhay.
- Mazur V., Didur I., Myalkovsky R., Pantsyрева H., Telekalo N., Tkach O. (2020). The productivity of intensive pea varieties depending on the seeds treatment and foliar fertilizing under conditions of right-bank forest-steppe Ukraine. *Ukrainian Journal of Ecology*, 10(1), 101–105.
- Mazur V.A., Didur I.M., Pantsyрева H.V., Telekalo N.V. (2018). Energy-economic efficiency of growth of grain-crop cultures in conditions of Right-Bank Forest-Steppe of Ukraine. *Ukrainian Journal of Ecology*, 8(4), 26–33.
- Mazur V.A., Mazur K.V., Pantsyрева H.V. (2019). Influence of the technological aspects growing on quality composition of seed white lupine (*Lupinus albus* L.) in the Forest Steppe of Ukraine. *Ukrainian Journal of Ecology*, 9, 50–55.
- Mazur V.A., Mazur K.V., Pantsyрева H.V., Alekseev O.O. (2018). Ecological and economic evaluation of varietal resources *Lupinus albus* L. in Ukraine. *Ukrainian Journal of Ecology*, 8, 148–153.
- Mazur V.A., Pantsyрева H.V., Mazur K.V., Didur I.M. (2019). Influence of the assimilation apparatus and productivity of white lupine plants. *Agronomy Research*, 17(X), 206–209. DOI <https://doi.org/10.15159/AR.19.024>.
- Mazur V.A., Pantsyрева H.V., Mazur K.V., Myalkovsky R.O., Alekseev O.O. (2020). Agroecological prospects of using corn hybrids for biogas production. *Agronomy Research* 18(1), 177–182.
- Mazur, V. A., & Pantsyрева, H. V. (2017). Vplyv tekhnolohichnykh pryiomiv vyroshchuvannya na urozhainist i yakist zerna liupynu biloho v umovakh Pravoberezhnoho Lisostepu. *Silske hospodarstvo i lisivnytstvo*, 7, 27–36.
- Mazur, V. A., Myalkovsky, R.O., Mazur, K. V., Pantsyрева, H. V., Alekseev, O.O. (2019). Influence of the Photosynthetic Productivity and Seed Productivity of White Lupine Plants. *Ukrainian Journal of Ecology*, 9(4), 665–670.
- Mazur, V.A., Branitskyi, Y.Y., Pantsyрева, H.V. (2020). Bioenergy and economic efficiency technological methods growing of switchgrass. *Ukrainian Journal of Ecology*, 10(2), 8–15.
- Mazur, V.A., Didur, I.M., Pantsyрева, H.V., & Telekalo, N.V. (2018). Energy-economic efficiency of grain-crop cultures in the conditions of the right-bank Forest-Steppe of Ukraine. *Ukrainian J Ecol*, 8(4), 26–33.
- Mazur, V.A., Mazur, K.V., Pantsyрева, H.V., Alekseev, O.O. (2018). Ecological and economic evaluation of varietal resources *Lupinus albus* L. in Ukraine. *Ukrainian Journal of Ecology*, 8(4), 148–153.
- Melnychuk, T., Patyka, V. (2011). Microbial preparations bioorganic farming system. *Proceed. Of Third All-Ukrainian Congress of Ecologists, Vinnytsya*, 2, 423–426.
- Metodologiya i praktyka vykorystannya mikrobnix preparativ u texnologiyax vyroshhuvannya silskogospodarskyx kultur [Methodology and practice of microbial drugs use in crop growing technologies]. (2011). V. V. Volkogon, A. S. Zaryshnyak, I. V. Grynyk (Eds.). Kyiv: Agrarna nauka.
- Mohamed Z., El-Sayed S., Radwan T., El-Wahab G. (2009). Potency evaluation of *Serratiamarcescens* and *Pseudomonas fluorescens* as biocontrol agents for root-knot nematodes in Egypt. *Journal of Applied Sciences Research*, 4(1), 93–102.
- Monarkh V.V., Pantsyрева, H.V. (2019). Stages of the Environmental Risk Assessment. *Ukrainian Journal of Ecology*, 9(4), 484–492. DOI: 10.15421/2019_779
- Muir J.P., Sanderson, M.A., Ocumpaugh, W.R. (2001). Biomass production of Alamo switchgrass in response to nitrogen, phosphorus, and row spacing. *Agron J.*, 93, 896–901.
- Osoro N., Kawaka F., Naluyange V. (2014). Effects of water hyacinth (*Eichhornia crassipes* [mart.] solms) compost on growth and yield of common beans (*Phaseolus vulgaris*) in Lake Victoria Basin. *European International Journal of Science and Technology*, 3(7), 173–186.
- Oliveira, I.A., Vasconcellos, M.J., Seldin, L., Paiva, E., Vargas, M.A., de Sá, N.M.H. (2000). Random Amplified Polymorphic DNA analysis of effective *Rhizobium* sp. associated with beans cultivated in Brazilian cerrado soils. *Braz J Microbiol*, 31, 39–44.
- Ovcharuk V.I., Mulyarchuk O.I., Myalkovsky R.O., Bezikonnyi P.V., Kravchenko V.S., Klymoych N.M. (2019). Parameters of beet plants. *Bulletin of the Uman National University of Horticulture*, 1, 70–75.
- Palamarchuk V., Honcharuk I., Honcharuk T., Telekalo N. (2018). Effect of the elements of corn cultivation the technology on bioethanol production under conditions of the rightbank forest-steppe of Ukraine. *Ukrainian Journal of Ecology*, 8(3), 47–53.
- Pantsyрева H.V. (2018). Research on varietal resources of herbaceous species of *Paeonia* L. in Ukraine. *Scientific Bulletin of the NLTU of Ukraine*, 28(8), 74–78. <https://doi.org/10.15421/40280815>
- Pantsyрева H.V., Myalkovsky R.O., Yasinetska I.A., Prokopchuk V.M. (2020). Productivity and economical appraisal of growing raspberry according to substrate for mulching under the conditions of podilia area in Ukraine. *Ukrainian Journal of Ecology*, 10(1), 210–214.
- Pantsyрева, H.V., Mykoliuk, O.O., & Semchuk, V.V. (2019). Suchasnyi stan kolektsii pivonii na bazi botanichnoho sadu "Podillia" Vinnytskoho natsionalnoho ahrarnoho universytetu. *Scientific Bulletin of UNFU*, 29(8), 46–50. <https://doi.org/10.36930/40290806>
- Pantsyрева, H.V. (2019). Morphological and ecological-biological evaluation of the decorative species of the genus *Lupinus* L.. *Ukrainian Journal of Ecology*, 9(3), 74–77.
- Pantsyрева, H.V. (2019). Technological aspects of biogas production from organic raw materials. *Bulletin of KhNTUSG them. P. Vasilenko. Kharkiv*.
- Pruvot E.B., Claeys-Kulik A.L., Estermann T. (2015). Designing strategies for efficient funding of universities in Europe. *DEFINE Project Paper*. Brussels: European University Association.

- Puyu V.L., Bakhmat M.I., Rykhivskiy I.P., Shcherbatiuk N.V. (2019). Optimization of Conveyor Production of Green Fodder. *World Science*, 7(47), 1. doi: 10.31435/rsglobal_ws/31072019/6587.
- Rai R.K., Tripathi N., Gautam D., & Singh P. (2017). Exogenous application of ethrel and gibberellic acid stimulates physiological growth of late planted sugarcane with short growth period in subtropical India. *Journal of Plant Growth Regulation*, 36(2), 472-486.
- Razanov S.F., Tkachuk O.P., Mazur V.A., Didur I.M. (2018). Effect of bean perennial plants growing on soil heavy metal concentrations. *Ukrainian Journal of Ecology*, 8(2), 294-300 doi: 10.15421/2018_341.
- Rohach V.V., Rohach T.I., Kylynyk A.M., Polyvanyi S.V., Bayurko N.V., Nikitchenko L.O., Tkachuk O.O., Shevchuk O.A., Hudzevych L.S., Levchuk N.V. (2020). The influence of synthetic growth promoters on morphophysiological characteristics and biological productivity of potato culture. *Modern Phytomorphology*, 14, 111-114.
- Shatilov I. S., Chudnovskiy A. F. (1980). *Agrophysical, Agrometeorological and Agrotechnical Crop Programming Fundamentals: ACS of TP Principles in Land Cultivation*. Leningrad: Gidrometeoizdat.
- Solona O., Kupchuk I. (2020). Dynamic synchronization of vibration exciters of the three-mass vibration mill. *Przegląd Elektrotechniczny*, 96(3), 161-165. DOI: 10.15199/48.2020.03.35
- Varchenko O., Krysanov D., Shubravska O., Khakhula L., Gavryk O., Byba V., Honcharuk I. (2020). Supply Chain Strategy in Modernization of State Support Instruments for Small Farms in Ukraine. *International Journal of Supply Chain Management*, 9(1), 536-543
- Vdovenko S.A., Prokopchuk V.M., Palamarchuk I.I., Pansyryeva H.V. (2018). Effectiveness of the application of soil milling in the growing of the squash (*Cucurbita pepo* var. *giraumontia*) in the right-bank forest stepp of Ukraine. *Ukrainian Journal of Ecology*, 8(4), 1-8.
- Vdovenko, S.A., Pansyryeva, G.V., Palamarchuk, I.I., & Lytvyniuk, H.V. (2018). Symbiotic potential of snap beans (*Phaseolus vulgaris* L.) depending on biological products in agrocoenosis of the right-bank forest-steppe of Ukraine. *Ukrainian J Ecol*, 8(3), 270-274.
- Vdovenko, S.A., Prokopchuk, V.M., Palamarchuk, I.I., & Pansyryeva, H.V. (2018). Effectiveness of the application of soil milling in the growing of the squash (*Cucurbita pepo* var. *giraumontia*) in the right-bank forest stepp of Ukraine. *Ukrainian J Ecol*, 8(4), 1-5.
- Wolters D., Beste A. (2000). Biomasse – umweltfreundlicher Energieträger? *Ökologie und Landbau*, 116(4), 12-14.
- Yanovych, V., Honcharuk, T., Honcharuk, I. & Kovalova, K. (2018). Engineering management of vibrating machines for targeted mechanical activation of premix components. *INMATEH - Agricultural Engineering*, 54(1), 25-32.
- Yhurber J.A. (1958). Inhibitory effect of gibberellins on nodulation in dwarf beans, *Phaseolus vulgaris*. *Nature*, 181, 1082-1083.
- Yowling W.A., Buirchell B.J., Tarta M.E. (1998). *Lupin. Lupinus L.*, Promoting the conservation and use of underutilized and neglected crops. Institute of Plant Iyenetis and crop Plant Research, yatersleben. International Plant Iyenetis Resources Institute. Rome, 112-114.
- Zhao, H., Cao, H., Ming-Zhen, P., Sun, Y., & Liu, T. (2017). The role of plant growth regulators in a plant aphid parasitoid tritrophic system. *Journal of Plant Growth Regulation*, 36(4), 868-876.

Citation:

Didur, I., Chynchyk, O., Pansyryeva, H., Olifirovych, S., Olifirovych, V., Tkachuk, O. (2021). Effect of fertilizers for *Phaseolus vulgaris* L. productivity in Western Forest-Steppe of Ukraine. *Ukrainian Journal of Ecology*, 11 (1), 419-424.



This work is licensed under a Creative Commons Attribution 4.0. License
