

ORIGINAL ARTICLE

New records of holothurian-associated pea crab *Pinnixa tumida* Stimpson, 1858 (Crustacea: Decapoda: Pinnotheridae) from the Russian coastal waters of the Sea of Japan

I. Marin

*A.N. Severtzov Institute of Ecology and Evolution of RAS, Moscow, 119071, Russia
Biological Department, Altai State University, Barnaul, 656049, Russia*

*E-mail: coralliodecapoda@mail.ru

Received: 24.10.2018. Accepted: 25.11.2018

Pea crab *Pinnixa tumida* Stimpson, 1858 associated with burrowing holothurian *Paracaudina chilensis* (Müller, 1850) is recorded from the Peter the Great Bay and Posjeta Bay of the Sea of Japan. The new locality where the species was found is situated about 130 km north-east from the previous record and presents the most northern distribution of the species in the Northern Pacific. The distribution of the species in other areas is discussed in the paper.

Keywords: Crustacea; Decapoda; Brachyura; Pinnotheridae; *Pinnixa tumida*; *Paracaudina*; pea crab; symbionts; new records; Sea of Japan

Introduction

The decapod fauna of Russian coastal waters of the Sea of Japan presently includes about 70 species from 22 families (Vinogradov 1950; Marin 2010, 2013a, c, 2017b; Marin & Kornienko 2014) including numerous species recently recorded from the area (Marin et al. 2011, 2013a–c, 2013; Marin 2013a–d, 2014, 2016a, b, 2017a, b, 2018a; Anker et al., 2016) as well as some species firstly found in other regions of the northern Pacific (Marin et al. 2015; Komai et al. 2017; Marin, 2018b) indicating that brachyuran and decapod fauna and the species distribution are still known incompletely.

Three species of pea crab genus *Pinnixa* White, 1846 (Crustacea: Decapoda: Pinnotheridae) have been recorded for the fauna of Russia, mostly from the western part of the Sea of Japan: *Pinnixa tumida* Stimpson, 1858, *P. rathbuni* Sakai, 1934 and *P. banzu* Komai, Nishi & Taru, 2014 (Vinogradov, 1950; Rybakov, 1986; Vasilenko, 1990; Sinelnikov & Marin, 2012; Marin, 2013a, b, 2015, 2016a; Marin & Kornienko, 2014). These species clearly differ morphologically as well as by ecological features. *P. tumida* was found in Posyet Bay in the intestine of the burrowing holothurian *Paracaudina chilensis* (Müller, 1850) (= *Caudina ransonnetti* Marenzeller von, 1881, a junior synonym) (Echinodermata: Holothuroidea: Caudinidae), *P. rathbuni* is living in burrows of the spoon worm *Urechis unicinctus* (von Drasche, 1881) (Annelida: Echiura: Urechidae) and *P. banzu* is known as inhabitant of the tubes of polychaete *Chaetopterus cautus* Marenzeller, 1879 (Annelida: Polychaeta: Chaetopteridae).

During an intensive study of decapod biodiversity along the Russian coast of the Sea of Japan, the species was found again from the northern part of the Peter the Great Bay that significantly increases the area of the species to the north.

Material and methods

Fieldwork and sampling were carried out along the Russian coasts of the Sea of Japan, in Peter the Great Bay (September 2017–2018) and Posyet Bay (July 2016). The material was collected from subtidal waters to the depth of 20 meters by snorkeling and SCUBA diving. Coloration of living specimens and their external morphology was photographed using digital camera Canon G16. All specimens were preserved in 70% ethanol for further DNA analysis. Carapace width (cw, mm), the maximal length between lateral margins of carapace, is used as standard measurement. Species names and modern taxonomic position are given according to Ng *et al.* (2008), Davie *et al.* (2015), the international database WoRMS (World Register of Marine Species) and Marine Species Identification Portal. The examined material is deposited in the personal author collection (LEMMI) deposited in A.N. Severtzov Institute of Ecology and Evolution of Russian Academy of Sciences, Moscow.

Results and discussion

Order Decapoda Latreille, 1803

Family Pinnotheridae De Haan, 1833

Genus *Pinnixa* White, 1846

Pinnixa tumida Stimpson, 1858

(Fig. 1)

Pinnixa tumida Stimpson, 1858: 108 [type locality: Hakodate Bay, Japan].

Material Examined. Sea of Japan, Russia, Posyeta Bay – 2 ovigerous females, 5 males (LEMMI), Vitjaz Bay, 42°36'05.0"N, 131°11'01.2"E, muddy sand bottom, on *Zostera* beds, from holothurian burrowing in the sand, depth 2–3 m, snorkeling, coll. I. Marin & A. Mayorova, 16 June 2016; the Peter the Great Bay – 1 ovigerous female, 1 male (LEMMI), Astafieva Bay, 42°36'52.2"N 131°12'01.1"E, depth 1–1.5 m, clear sand bottom, from burrowing holothurian, hand sampling, coll. Marin I., August 2018; 2 ovigerous females, 2 males (LEMMI), Vostok Bay, 42°53'10.0"N, 132°43'42.9"E, sandy bottom, from holothurian burrowing in the sand, 10–15 m., SCUBA, coll. K. Dudka, August 2018.

Remarks. In contrast to the other representatives of the genus *Pinnixa* occurring in the Russian waters, *P. tumida* has a tooth in the middle of the cutting edge of the movable finger (dactylus) of the chela of pereopod I (see Fig. 1c), while in *P. rathbuni* and *P. banzu* it is situated at the base of the immovable finger (polex) of the chela. The carapace width of sexually mature *P. tumida* is with the convex and strong calcified that also clearly differ the species from the congeners in the Sea of Japan.

Coloration. One of brightly colored species within the genus. Dorsal surface of carapace and chela colored with brown and violet spots; setae on appendages deeply brown (see Fig. 1b, c). The reddish brown pigment in the carapace of these crabs, frequently retained in preserved specimens, is thought to be derived from carotenoids originally ingested by the host (Hopkins & Scanland, 1964; after Zmarzly, 1992).

Size. The largest female has cw. 14 mm, male has cw. 10 mm; the females are larger than the males.




Figure 1. The pea crab *Pinnixa tumida* Stimpson, 1858 and its host – a burrowing holothurian *Paracaudina chillensis* (Müller, 1850): *a*, *b* – general view; *c* – frontal view.

Habitat and ecology. Subtidal species. The crab lives in the intestine of the burrowing holothurian *P. chillensis* (Konishi, 1977; Vassilenko, 1990; Takeda et al., 1997) probably representing an undescribed species of "*P. chillensis*" species complex)

(=*Caudina ransonnetti* Marenzeller von, 1881, a junior synonym). The species is mostly ecologically similar to *P. barnharti* Rathbun, 1918 living in the body cavity of the burrowing holothurian *Molpadia arenicola* (Stimpson, 1858), as well as holothurians of the genera *Chiridota* Eschscholtz, 1829 and *Caudina* Stimpson, 1853 along the Pacific coast of North America (Schmitt et al., 1973; Morris et al., 1980; Zmarzly, 1992).

Distribution. In Russian waters, *P. tumida* presently known from Posyeta Bay and the Peter the Great Bay of the Sea of Japan. In the Sea of Japan, this species was also found at the coasts of islands of Japan, Korean Peninsula, and in the Yellow Sea along the eastern coast of China (Kim, 1973; Konishi, 1977; Vassilenko, 1990; Dai & Yang, 1991; Komai et al., 1993).

Acknowledgements

Author is very thankful to A. Mayorova and K. Dudka for the help during the sampling in Posyeta Bay and the Peter the Great Bay, respectively.

References

- Anker, A., Marin, I., Pachelles, P.P.G. & Komai, T. (2016). Redescription of *Alpheus vladivostokiensis* (Vinogradov, 1950), a large and conspicuous snapping shrimp from the northern Sea of Japan (Decapoda: Caridea: Alpheidae). *Zootaxa*, 4127(1), 171–184.
- Davie, P.J., Guinot, D. & Ng, P.K. (2015). Systematics and classification of Brachyura. In: P. Castro, P.J.F. Davie, D. Guinot, F.R. Schram & von Vaupel Klein, J.C. (eds). *Treatise on Zoology – Anatomy, Taxonomy, Biology. The Crustacea*, 9, Part C (2 vol.). Leiden, Brill, 1049–1130.
- Dai, A. & Yang, S. (1991). *Crabs of the China Seas*. Beijing: China Ocean Press and Berlin: Springer-Verlag. 345 pp.
- Hopkins, T.S. & Scanland, T.B. (1964) The host relations of a pinnotherid crab, *Opisthopus transversus* Rathbun (Crustacea: Decapoda). *Bulletin of the Southern California Academy of Sciences*, 63, 175–180.
- Kim, H.S. (1973). *A Catalogue of Anomura and Brachyura from Korea*. Illustrated Encyclopedia of Fauna and Flora of Korea, Seoul: Samwha Publishing Co., 14, 1–694.
- Komai, T., Maruyama, S. & Konishi, K. (1992). A list of decapod crustaceans from Hokkaido, northern Japan. *Researches on Crustacea*, 21, 189–205.
- Komai, T., Marin, I. & Kakui, K. (2017). Rediscovery and redescription of the abyssal squat lobster *Munidopsis petalorhyncha* Baba, 2005 (Crustacea: Decapoda: Munidopsidae) from the Northwest Pacific. *Zootaxa*, 4226(1), 93–102.
- Konishi, K. (1977). On three species of the pinnotherid crabs from Hokkaido, Japan. *Journal of the Faculty of Science, Hokkaido University: Series 6. Zoology*, 20(4), 605–610.
- Marin, I. (2010). Redescription of the alpheid shrimp *Betaeus levifrons* Vinogradov, 1950 (Crustacea, Decapoda, Alpheidae) from Peter the Great Bay, Russian coast of the Sea of Japan. *Zootaxa*, 2613, 51–60.
- Marin, I.N. (2013a). Infraorder Brachyura. In: Sirenko, B.I. (ed). *Check-list of species of free-living invertebrates of the Russian Far Eastern Seas*. Zoological Institute RAS, St. Petersburg. 250 pp.
- Marin, I.N. (2013b). New data on the distribution of hairy-clawed shore crabs of the genus *Hemigrapsus* (Decapoda: Varunidae) along the Russian mainland coast of the Sea of Japan. *Russian Journal of Marine Biology*, 39(4), 301–305.
- Marin, I.N. (2013c). *Atlas of Decapod Crustaceans of Russia*. KMK Scientific Press, Moscow, 145 pp. (in Russian, with English abstract).
- Marin, I. (2013d). A new species of callianassid ghost shrimp of the genus *Nihonotrypaea* Manning & Tamaki, 1998 (Crustacea, Decapoda, Axiidea, Callianasidae) from southern Posyeta Bay of the Russian coast of the Sea of Japan. *Zootaxa*, 3694(5), 434–444.
- Marin, I.N. (2014). Finding of the pea crab *Pinnaxodes mutuensis* Sakai, 1939 (Crustacea: Decapoda: Pinnotheridae) in an unusual host in Busse Lagoon, southern Sakhalin. *Russian Journal of Marine Biology*, 40(6), 486–489.
- Marin, I.N. (2016a). The species composition and ecological features of pea crabs of the genus *Pinnixa* White, 1846 (Brachyura: Pinnotheridae) in Peter the Great Bay, the Sea of Japan. *Russian Journal of Marine Biology*, 42(2), 139–145.
- Marin, I. (2016b). Notes on holotypes of hermit crabs (Decapoda: Anomura: Paguroidea) deposited in the collection of Zoological Museum of RAS (Saint-Petersburg, Russia) with remarks on hermit crab diversity along the Russian coast of the Sea of Japan. *Zootaxa*, 4105(2): 171–180.
- Marin, I. (2017a). *Athanas alpheusophilus* sp. nov. (Decapoda: Alpheidae) – a new *Alpheus*-associated shrimp from the Russian coast of the Sea of Japan. *Zootaxa*, 4324(12), 50–62.
- Marin I.N. (2017b). Estuarine crabs of the Okhotsk Sea and the Sea of Japan: rare species and new records. *Ukrainian Journal of Ecology*, 7(4), 444–452.
- Marin, I. (2018a). On the taxonomic status of amphidromous shrimp *Paraty borealis* Volk, 1938 (Crustacea: Decapoda: Atyidae) from the south of the Russian Far East. *Zootaxa*, 4444(2), 154–162.
- Marin, I. (2018b). Deep water decapod crustaceans (Crustacea: Decapoda) collected by SochoBio 2015 Expedition from bathyal and abyssal waters of the Sea of Okhotsk and adjacent NW Pacific with the re-description of *Calocarides okhotskensis* Sakai, 2011 (Axiidae). Deep Sea Research Part II: Topical Studies in Oceanography, 154, 330–341. <https://doi.org/10.1016/j.dsr2.2018.04.007>

- Marin, I., Mayorova, A. & Korn, O. (2018). Cryptic diversity of the rocky crab genus *Glebocarcinus* Nations, 1975 (Crustacea: Decapoda: Cancridae): description of a new species from Russian coastal waters of the Sea of Japan based on morphology, DNA and distribution. *Zootaxa*, 4415(3), 473–497.
- Marin, I.N. & Kornienko E.S. (2014). The list of Decapoda species from Vostok Bay Sea of Japan. *Biodiversity and Environment of Far East Reserves*, 2, 49–71 (in Russian).
- Marin, I.N., Korn, O.M. & Kornienko, E.S. (2011). Symbiotic crab *Sestrostoma balssi* (Shen, 1932) (Varunidae: Gaeticinae) from Vostok Bay, Sea of Japan: a new species for the fauna of Russia. *Russian Journal of Marine Biology*, 37(6), 509–511.
- Marin, I.N., Korn, O.M. & Kornienko, E.S. (2012a). The caridean shrimp *Lysmata vittata* (Stimpson, 1860) (Decapoda: Hippolytidae): a new species for the fauna of Russia. *Russian Journal of Marine Biology*, 38(4), 359–363.
- Marin, I.N., Korn, O.M. & Kornienko, E.S. (2012b). On the presence of the caridean shrimp *Lysmata vittata* (Decapoda, Hippolytidae) in the Russian waters of the Sea of Japan. *Crustaceana*, 85(1), 123–124.
- Marin, I.N., Korn, O.M. & Kornienko, E.S. (2012c). Hermit Crabs *Pagurus parvispina* Komai, 1997 and *Discorsopagurus macLaughlinae* Komai, 1995 (Decapoda: Paguridae): New Records for Russian Waters of the Sea of Japan. *Russian Journal of Marine Biology*, 38(3): 275–278. DOI: 10.1134/S1063074012030091
- Marin, I.N., Korn, O.M. & Kornienko, E.S. (2013). *Upogebia yokoyai* Makarov, 1938 (Decapoda: Upogebiidae): a gebiid shrimp species new to the fauna of the Sea of Japan. *Biologiya Morya*, 39(3), 221–226 (in Russian).
- Marin, I.N., Ng, P.K.L. & Anosov, S.E. (2015). A new record of the Japanese Longhorn decorator crab *Chorilia japonica* (Miers, 1879) (Crustacea, Brachyura, Epialtidae) in the western part of Bering Sea – the first record of the species and genus for Russian waters. *Arthropoda Selecta*, 24(2), 185–188.
- Ng, P.K.L., Guinot, D. & Davie, P.J.F. (2008). Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. *Raffles Bulletin of Zoology*, 17, 1–286.
- Schmitt, W.L., McCain, J.C. & Davidson, E.S. (1973). *Crustaceorum Catalogus. Pt. 3. Decapoda I, Brachyura I, Fam. Pinnotheridae*. Junk, the Hauge. 160 pp.
- Sinel'nikov, S.Yu. & Marin, I.N. (2012). Inventarizatsiya fauny desyatinogih rakoobraznyh Rossii kak odnogo iz osnovnyh morskikh biologicheskikh resursov. *Reports of Far Eastern State Technical Fisheries University*, 26, 24–27 (in Russian).
- Takeda S., Tamura S. & Washio M. (1997). Relationship between the pea crab *Pinnixa tumida* and its endobenthic holothurian host *Paracaudina chilensis*. *Marine Ecology Progress Series*, 149, 143–154.
- Vassilenko, S.V. (1990). On taxonomy and ecology of commensal crabs of family Pinnotheridae (Crustacea, Decapoda, Brachyura) of the Sea Okhotsk and northern part of the Japan Sea. *Proceedings of the Zoological Institute of RAS*, 218, 75–95.
- Vinogradov, L.G. (1950). Identification keys for shrimps, prawns and crabs from the Far East. *Izvestiya TINRO*, 33, 179–358 (in Russian).
- WoRMS Editorial Board (2017). World Register of Marine Species. Available from: <http://www.marinespecies.org> (accessed 21 October 2017).
- Zmarzly, D.L. (1992). Taxonomic review of pea crabs in the genus *Pinnixa* (Decapoda: Brachyura: Pinnotheridae) occurring on the California shelf, with descriptions of two species. *Journal of Crustacean Biology*, 12(4), 677–713.

Citation:

Marin, I. (2018). New records of holothurian-associated pea crab *Pinnixa tumida* Stimpson, 1858 (Crustacea: Decapoda: Pinnotheridae) from the Russian coastal waters of the Sea of Japan. *Ukrainian Journal of Ecology*, 8(4), 307–310



This work is licensed under a Creative Commons Attribution 4.0. License
